

DOMAIN OF DIFFERENCE MATRIX OF ORDER ONE IN SOME SPACES OF DOUBLE SEQUENCES

SERKAN DEMİRİZ AND OSMAN DUYAR

ABSTRACT. In this study, we define the spaces $\mathcal{M}_u(\Delta)$, $\mathcal{C}_p(\Delta)$, $\mathcal{C}_{0p}(\Delta)$, $\mathcal{C}_r(\Delta)$ and $\mathcal{L}_q(\Delta)$ of double sequences whose difference transforms are bounded, convergent in the Pringsheim's sense, null in the Pringsheim's sense, both convergent in the Pringsheim's sense and bounded, regularly convergent and absolutely q -summable, respectively, and also examine some inclusion relations related to those sequence spaces. Furthermore, we show that these sequence spaces are Banach spaces. We determine the alpha-dual of the space $\mathcal{M}_u(\Delta)$ and the $\beta(v)$ -dual of the space $\mathcal{C}_\eta(\Delta)$ of double sequences, where $v, \eta \in \{p, bp, r\}$. Finally, we characterize the classes $(\mu : \mathcal{C}_v(\Delta))$ for $v \in \{p, bp, r\}$ of four dimensional matrix transformations, where μ is any given space of double sequences.

1. INTRODUCTION

By ω and Ω , we denote the sets of all real valued single and double sequences which are the vector spaces with coordinatewise addition and scalar multiplication. Any vector subspaces of ω and Ω are called as the *single sequence space* and *double sequence space*, respectively. By $\mathcal{M}_u$, we denote the space of all bounded double sequences, that is

$$\mathcal{M}_u := \left\{ x = (x_{mn}) \in \Omega : \|x\|_\infty = \sup_{m,n \in \mathbb{N}} |x_{mn}| < \infty \right\}$$

which is a Banach space with the norm $\|x\|_\infty$; where $\mathbb{N}$ denotes the set of all positive integers. Consider a sequence $x = (x_{mn}) \in \Omega$. If for every $\varepsilon > 0$ there exists $n_0 = n_0(\varepsilon) \in \mathbb{N}$ and $l \in \mathbb{R}$ such that $|x_{mn} - l| < \varepsilon$ for all $m, n > n_0$ then we call that the double sequence x is *convergent* in the *Pringsheim's sense* to the limit l and write $p - \lim x_{mn} = l$; where $\mathbb{R}$ denotes the real field. By $\mathcal{C}_p$, we denote the space of all convergent double sequences in the Pringsheim's sense. It is well-known that there are such sequences in the space $\mathcal{C}_p$ but not in the space $\mathcal{M}_u$. Indeed following Boos [1], if we define the sequence $x = (x_{mn})$ by

$$x_{mn} := \begin{cases} n & , \quad m = 1, n \in \mathbb{N}, \\ 0 & , \quad m \geq 2, n \in \mathbb{N}, \end{cases}$$

then it is trivial that $x \in \mathcal{C}_p \setminus \mathcal{M}_u$, since $p - \lim x_{mn} = 0$ but $\|x\|_\infty = \infty$. So, we can consider the space $\mathcal{C}_{bp}$ of the double sequences which are both convergent in the Pringsheim's sense and bounded, i.e., $\mathcal{C}_{bp} = \mathcal{C}_p \cap \mathcal{M}_u$. A sequence in the space $\mathcal{C}_p$ is said to be *regularly convergent* if it is a single convergent sequence with respect to each index and denote the set of all such sequences by $\mathcal{C}_r$. Also by $\mathcal{C}_{bp0}$ and $\mathcal{C}_{r0}$, we denote the spaces of all double sequences converging to 0 contained in the sequence spaces $\mathcal{C}_{bp}$ and $\mathcal{C}_r$, respectively. Móricz [2] proved that $\mathcal{C}_{bp}$, $\mathcal{C}_{bp0}$, $\mathcal{C}_r$ and $\mathcal{C}_{r0}$ are Banach spaces with the norm $\|\cdot\|_\infty$.

Let us consider the isomorphism T defined by Zeltser [3] as

$$(1.1) \quad \begin{aligned} T : \Omega &\rightarrow \omega \\ x &\mapsto z = (z_i) := (x_{\varphi^{-1}(i)}), \end{aligned}$$

where $\varphi : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ is a bijection defined by

2000 *Mathematics Subject Classification.* 40C05, 46A45 .

Key words and phrases. Double sequence space, Matrix transformations,

$$\begin{aligned}
\varphi[(1,1)] &= 1, \\
\varphi[(1,2)] &= 2, \varphi[(2,2)] = 3, \varphi[(2,1)] = 4, \\
&\cdot \\
&\cdot \\
&\cdot \\
\varphi[(1,n)] &= (n-1)^2 + 1, \varphi[(2,n)] = (n-1)^2 + 2, \dots, \\
\varphi[(n,n)] &= (n-1)^2 + n, \varphi[(n,n-1)] = n^2 - n + 2, \dots, \varphi[(n,1)] = n^2, \\
&\cdot \\
&\cdot \\
&\cdot
\end{aligned}$$

Let us consider a double sequence $x = (x_{mn})$ and define the sequence $s = (s_{mn})$ which will be used throughout via x by

$$(1.2) \quad s_{mn} := \sum_{i=0}^m \sum_{j=0}^n x_{ij}$$

for all $m, n \in \mathbb{N}$. For the sake of brevity, here and in what follows, we abbreviate the summation $\sum_{i=0}^{\infty} \sum_{j=0}^{\infty}$ by $\sum_{i,j}$ and we use this abbreviation with other letters. Let λ be a space of a double sequences, converging with respect to some linear convergence rule $v - \lim : \lambda \rightarrow \mathbb{R}$. The sum of a double series $\sum_{i,j} x_{ij}$ with respect to this rule is defined by $v - \sum_{i,j} x_{ij} = v - \lim_{m,n \rightarrow \infty} s_{mn}$. Let λ, μ be two spaces of double sequences, converging with respect to the linear convergence rules $v_1 - \lim$ and $v_2 - \lim$, respectively, and $A = (a_{mnkl})$ also be a four dimensional infinite matrix over the real or complex field.

The α -dual λ^α , $\beta(v)$ -dual $\lambda^{\beta(v)}$ with respect to the v -convergence for $v \in \{p, bp, r\}$ and the γ -dual λ^γ of a double sequence space λ are respectively defined by

$$\begin{aligned}
\lambda^\alpha &:= \left\{ (a_{ij}) \in \Omega : \sum_{i,j} |a_{ij} x_{ij}| < \infty \text{ for all } (x_{ij}) \in \lambda \right\}, \\
\lambda^{\beta(v)} &:= \left\{ (a_{ij}) \in \Omega : v - \sum_{i,j} a_{ij} x_{ij} \text{ exists for all } (x_{ij}) \in \lambda \right\}, \\
\lambda^\gamma &:= \left\{ (a_{ij}) \in \Omega : \sup_{k,l \in \mathbb{N}} \left| \sum_{i,j=1}^{k,l} a_{ij} x_{ij} \right| < \infty \text{ for all } (x_{ij}) \in \lambda \right\}.
\end{aligned}$$

It is easy to see for any two spaces λ, μ of double sequences that $\mu^\alpha \subset \lambda^\alpha$ whenever $\lambda \subset \mu$ and $\lambda^\alpha \subset \lambda^\gamma$. Additionally, it is known that the inclusion $\lambda^\alpha \subset \lambda^{\beta(v)}$ holds while the inclusion $\lambda^{\beta(v)} \subset \lambda^\gamma$ does not hold, since the v -convergence of the sequence of partial sums of a double series does not imply its boundedness.

The v -summability domain $\lambda_A^{(v)}$ of a four dimensional infinite matrix $A = (a_{mnkl})$ in a space λ of a double sequences is defined by

$$(1.3) \quad \lambda_A^{(v)} = \left\{ x = (x_{kl}) \in \Omega : Ax = \left(v - \sum_{k,l} a_{mnkl} x_{kl} \right)_{m,n \in \mathbb{N}} \text{ exists and is in } \lambda \right\}.$$

We say, with the notation (1.3), that A maps the space λ into the space μ if and only if Ax exists and is in μ for all $x \in \lambda$ and denote the set of all four dimensional matrices, transforming the space λ into the space μ , by $(\lambda : \mu)$. It is trivial that for any matrix $A \in (\lambda : \mu)$, $(a_{mnkl})_{k,l \in \mathbb{N}}$ is in the $\beta(v)$ -dual $\lambda^{\beta(v)}$ of the space λ for all $m, n \in \mathbb{N}$. An infinite matrix A is said to be $\mathcal{C}_v$ -conservative if $\mathcal{C}_v \subset (\mathcal{C}_v)_A$. Also by $(\lambda : \mu; p)$, we denote the class of all four dimensional matrices $A = (a_{mnkl})$ in the class $(\lambda : \mu)$ such that $v_2 - \lim Ax = v_1 - \lim x$ for all $x \in \lambda$.

Now, following Zeltser [4] we note the terminology for double sequence spaces. A locally convex double sequence space λ is called a *DK-space*, if all of the seminorms $r_{kl} : \lambda \rightarrow \mathbb{R}$, $x = (x_{kl}) \mapsto |x_{kl}|$ for all $k, l \in \mathbb{N}$ are continuous. A *DK-space* with a Fréchet topology is called an *FDK-space*. A normed *FDK-space* is called a *BDK-space*. We record that C_r endowed with the norm $\|\cdot\|_\infty : C_r \rightarrow \mathbb{R}$, $x = (x_{kl}) \mapsto \sup_{k,l \in \mathbb{N}} |x_{kl}|$ is a *BDK-space*.

Let us define the following sets of double sequences:

$$\begin{aligned}\mathcal{M}_u(t) &:= \left\{ (x_{mn}) \in \Omega : \sup_{m,n \in \mathbb{N}} |x_{mn}|^{t_{mn}} < \infty \right\}, \\ \mathcal{C}_p(t) &:= \left\{ (x_{mn}) \in \Omega : \exists l \in \mathbb{C} \ni p - \lim_{m,n \rightarrow \infty} |x_{mn} - l|^{t_{mn}} = 0 \right\}, \\ \mathcal{C}_{0p}(t) &:= \left\{ (x_{mn}) \in \Omega : p - \lim_{m,n \rightarrow \infty} |x_{mn}|^{t_{mn}} = 0 \right\}, \\ \mathcal{L}_u(t) &:= \left\{ (x_{mn}) \in \Omega : \sum_{m,n} |x_{mn}|^{t_{mn}} < \infty \right\}, \\ \mathcal{C}_{bp}(t) &:= \mathcal{C}_p(t) \cap \mathcal{M}_u(t) \quad \text{and} \quad \mathcal{C}_{0bp}(t) := \mathcal{C}_{0p}(t) \cap \mathcal{M}_u(t); \end{aligned}$$

where $t = (t_{mn})$ is the sequence of strictly positive reals t_{mn} for all $m, n \in \mathbb{N}$. In the case $t_{mn} = 1$ for all $m, n \in \mathbb{N}$; $\mathcal{M}_u(t), \mathcal{C}_p(t), \mathcal{C}_{0p}(t), \mathcal{L}_u(t), \mathcal{C}_{bp}(t)$ and $\mathcal{C}_{0bp}(t)$ reduce to the sets $\mathcal{M}_u, \mathcal{C}_p, \mathcal{C}_{0p}, \mathcal{L}_u, \mathcal{C}_{bp}$ and $\mathcal{C}_{0bp}$, respectively. Now, we can summarize the knowledge given in some document related to the double sequence spaces. Gökhan and Çolak [5, 6] have proved that $\mathcal{M}_u(t), \mathcal{C}_p(t)$ and $\mathcal{C}_{bp}(t)$ are complete paranormed spaces of double sequences and gave the alpha-, beta-, gamma-duals of the spaces $\mathcal{M}_u(t)$ and $\mathcal{C}_{bp}(t)$. Quite recently, in her PhD thesis, Zeltser [3] has essentially studied both the theory of topological double sequence spaces and the theory of summability of double sequences. Mursaleen and Edely [7] have introduced the statistical convergence and statistical Cauchy for double sequences, and gave the relation between statistically convergent and strongly Cesàro summable double sequences. Nextly, Mursaleen [8] and Mursaleen and Edely [9] have defined the almost strong regularity of matrices for double sequences and applied these matrices to establish a core theorem and introduced the M -core for double sequences and determined those four dimensional matrices transforming every bounded double sequence $x = (x_{jk})$ into one whose core is a subset of the M -core of x . More recently, Altay and Başar [10] have defined the spaces $\mathcal{BS}, \mathcal{BS}(t), \mathcal{CS}_{bp}, \mathcal{CS}_r$ and $\mathcal{BV}$ of double series whose sequence of partial sums are in the spaces $\mathcal{M}_u, \mathcal{M}_u(t), \mathcal{C}_p, \mathcal{C}_{bp}, \mathcal{C}_r$ and $\mathcal{L}_u$, respectively, and also examined some properties of those sequence spaces and determined the alpha-duals of the spaces $\mathcal{BS}, \mathcal{BV}, \mathcal{CS}_{bp}$ and the $\beta(v)$ -duals of the spaces $\mathcal{CS}_{bp}$ and $\mathcal{CS}_r$ of double series. Quite recently, Başar and Sever [11] have introduced the Banach space $\mathcal{L}_q$ of double sequences corresponding to the well-known space ℓ_q of absolutely q -summable single sequences and examine some properties of the space $\mathcal{L}_q$. Furthermore, they determine the $\beta(v)$ -dual of the space and establish that the alpha- and gamma-duals of the space $\mathcal{L}_q$ coincide with the $\beta(v)$ -dual; where

$$\begin{aligned}\mathcal{L}_q &:= \left\{ (x_{ij}) \in \Omega : \sum_{i,j} |x_{ij}|^q < \infty \right\}, \quad (1 \leq q < \infty), \\ \mathcal{CS}_v &:= \{ (x_{ij}) \in \Omega : (s_{mn}) \in \mathcal{C}_v \}. \end{aligned}$$

Here and after we assume that $v \in \{p, bp, r\}$.

The double difference matrix $\Delta = (\delta_{mnkl})$ of order one is defined by

$$\delta_{mnkl} := \begin{cases} (-1)^{m+n-k-l} & , \quad m-1 \leq k \leq m, \quad n-1 \leq l \leq n, \\ 0 & , \quad \text{otherwise} \end{cases}$$

for all $m, n, k, l \in \mathbb{N}$. Define the sequence $y = (y_{mn})$ as the Δ -transform of a sequence $x = (x_{mn})$, i.e.,

$$(1.4) \quad y_{mn} = (\Delta x)_{mn} = x_{mn} - x_{m,n-1} - x_{m-1,n} + x_{m-1,n-1}$$

for all $m, n \in \mathbb{N}$. Additionally, a direct calculation gives the inverse $\Delta^{-1} = S = (s_{mnkl})$ of the matrix Δ as follows:

$$s_{mnkl} := \begin{cases} 1 & , \quad 0 \leq k \leq m, \quad 0 \leq l \leq n, \\ 0 & , \quad \text{otherwise} \end{cases}$$

for all $m, n, k, l \in \mathbb{N}$.

In the present paper, we introduce the new double difference sequence spaces $\mathcal{M}_u(\Delta), \mathcal{C}_p(\Delta), \mathcal{C}_{0p}(\Delta)$ and $\mathcal{L}_q(\Delta)$, that is,

$$\begin{aligned} \mathcal{M}_u(\Delta) &:= \{(x_{mn}) \in \Omega : \sup_{m,n \in \mathbb{N}} |y_{mn}| < \infty\}, \\ \mathcal{C}_p(\Delta) &:= \{(x_{mn}) \in \Omega : \exists l \in \mathbb{C} \exists p - \lim_{m,n \rightarrow \infty} |y_{mn} - l| = 0\}, \\ \mathcal{C}_{0p}(\Delta) &:= \{(x_{mn}) \in \Omega : p - \lim_{m,n \rightarrow \infty} |y_{mn}| = 0\}, \\ \mathcal{L}_q(\Delta) &:= \left\{ (x_{ij}) \in \Omega : \sum_{m,n} |y_{mn}|^q < \infty \right\}, \quad (1 \leq q < \infty). \end{aligned}$$

By $\mathcal{C}_{bp}(\Delta)$ and $\mathcal{C}_r(\Delta)$, we denote the sets of all the Δ -transforms convergent and bounded, and the Δ -transforms regularly convergent double sequences. One can easily see that the spaces $\mathcal{M}_u(\Delta), \mathcal{C}_p(\Delta), \mathcal{C}_{0p}(\Delta), \mathcal{C}_{bp}(\Delta), \mathcal{C}_r(\Delta)$ and $\mathcal{L}_q(\Delta)$ are the domain of the double difference matrix Δ in the spaces $\mathcal{M}_u, \mathcal{C}_p, \mathcal{C}_{0p}, \mathcal{C}_{bp}, \mathcal{C}_r$ and $\mathcal{L}_q$, respectively.

2. SOME NEW DOUBLE DIFFERENCE SEQUENCE SPACES

In the present section, we deal with the sets $\mathcal{M}_u(\Delta), \mathcal{C}_p(\Delta), \mathcal{C}_{0p}(\Delta), \mathcal{C}_{bp}(\Delta), \mathcal{C}_r(\Delta)$ and $\mathcal{L}_q(\Delta)$ consisting of the double sequences whose Δ -transforms of order one are in the spaces $\mathcal{M}_u, \mathcal{C}_p, \mathcal{C}_{0p}, \mathcal{C}_{bp}, \mathcal{C}_r$ and $\mathcal{L}_q$, respectively.

Theorem 2.1. *The sets $\mathcal{M}_u(\Delta), \mathcal{C}_p(\Delta), \mathcal{C}_{0p}(\Delta), \mathcal{C}_{bp}(\Delta), \mathcal{C}_r(\Delta)$ and $\mathcal{L}_q(\Delta)$ are the linear spaces with the coordinatewise addition and scalar multiplication, and $\mathcal{M}_u(\Delta), \mathcal{C}_p(\Delta), \mathcal{C}_{0p}(\Delta), \mathcal{C}_{bp}(\Delta), \mathcal{C}_r(\Delta)$ and $\mathcal{L}_q(\Delta)$ are the Banach spaces with the norms*

$$(2.1) \quad \|x\|_{\mathcal{M}_u(\Delta)} = \sup_{m,n \in \mathbb{N}} |x_{mn} + x_{m-1,n-1} - x_{m,n-1} - x_{m-1,n}|,$$

$$(2.2) \quad \|x\|_{\mathcal{L}_q(\Delta)} = \left[\sum_{m,n} |x_{mn} + x_{m-1,n-1} - x_{m,n-1} - x_{m-1,n}|^q \right]^{1/q}, \quad (1 \leq q < \infty).$$

Proof. The first part of the theorem is a routine verification. So, we omit the detail.

Since the proof may be given for the spaces $\mathcal{M}_u(\Delta), \mathcal{C}_p(\Delta), \mathcal{C}_{0p}(\Delta), \mathcal{C}_{bp}(\Delta)$ and $\mathcal{C}_r(\Delta)$, to avoid the repetition of the similar statements, we prove the theorem only for the space $\mathcal{L}_q(\Delta)$.

It is obvious that $\|x\|_{\mathcal{L}_q(\Delta)} = \|y\|_q$, where $\|\cdot\|_q$ is the norm on the space $\mathcal{L}_q$. Let $x^{(r)} = \{x_{jk}^{(r)}\}$ be a Cauchy sequence in $\mathcal{L}_q(\Delta)$. Then, $\{y^{(r)}\}_{r \in \mathbb{N}}$ is a Cauchy sequence in $\mathcal{L}_q$, where $y^{(r)} = \{y_{mn}^{(r)}\}_{m,n=0}^\infty$ with

$$y_{mn}^{(r)} = x_{mn}^{(r)} + x_{m-1,n-1}^{(r)} - x_{m,n-1}^{(r)} - x_{m-1,n}^{(r)}$$

for all $m, n, r \in \mathbb{N}$. Then, for a given $\varepsilon > 0$, there is a positive integer $N = N(\varepsilon)$ such that

$$(2.3) \quad \|y^{(r)} - y^{(s)}\|_q = \left\{ \sum_{m,n} |y_{mn}^{(r)} - y_{mn}^{(s)}|^q \right\}^{1/q} < \varepsilon$$

for all $r, s > N$. Therefore $|y_{mn}^{(r)} - y_{mn}^{(s)}| < \varepsilon$, i.e. $\{y_{mn}^{(r)}\}_{r \in \mathbb{N}}$ is a Cauchy sequence in $\mathbb{C}$, and hence converges in $\mathbb{C}$. Say,

$$(2.4) \quad \lim_{r \rightarrow \infty} y_{mn}^{(r)} = y_{mn}.$$

Using these infinitely many limits, we define the sequence $y = (y_{mn})_{m,n=0}^\infty$. Then, we get by (2.4) that

$$(2.5) \quad \lim_{r \rightarrow \infty} \|y_{mn}^{(r)} - y_{mn}\|_{\mathcal{L}_q(\Delta)} = \lim_{r \rightarrow \infty} \left\{ \sum_{m,n} |y_{mn} - y_{mn}^{(r)}|^q \right\}^{1/q} = 0.$$

Now, we have to show that $y \in \mathcal{L}_q$. Since $y^{(r)} = \{y_{mn}^{(r)}\}_{m,n=0}^\infty \in \mathcal{L}_q$ and by (2.5)

$$\left\{ \sum_{m,n} |y_{mn}|^q \right\}^{1/q} \leq \left\{ \sum_{m,n} |y_{mn} - y_{mn}^{(r)}|^q \right\}^{1/q} + \left\{ \sum_{m,n} |y_{mn}^{(r)}|^q \right\}^{1/q} < \infty,$$

which shows that $y = (y_{mn})_{m,n=0}^\infty \in \mathcal{L}_q$, i.e. $x = (x_{jk}) \in \mathcal{L}_q(\Delta)$. Since $\{x^{(r)}\}_{r \in \mathbb{N}}$ was arbitrary Cauchy sequence in $\mathcal{L}_q(\Delta)$, the double difference sequence space $\mathcal{L}_q(\Delta)$ is complete. This completes the proof of the theorem. $\square$

Theorem 2.2. *The space $\lambda(\Delta)$ is linearly isomorphic to the space λ , where λ denotes any of the spaces $\mathcal{M}_u, \mathcal{C}_p, \mathcal{C}_{0p}, \mathcal{C}_{bp}, \mathcal{C}_r$ and $\mathcal{L}_q$.*

Proof. We show here that $\mathcal{M}_u(\Delta)$ is linearly isomorphic to $\mathcal{M}_u$. Consider the transformation T from $\mathcal{M}_u(\Delta)$ to $\mathcal{M}_u$ defined by $x = (x_{jk}) \mapsto y = (y_{mn})$. Then, clearly T is linear and injective. Now, define the sequence $x = (x_{jk})$ by

$$(2.6) \quad x_{jk} = \sum_{m=0}^j \sum_{n=0}^k y_{mn}$$

for all $j, k \in \mathbb{N}$. Suppose that $y \in \mathcal{M}_u$. Then, since

$$\begin{aligned} \|x\|_{\mathcal{M}_u(\Delta)} &= \sup_{j,k \in \mathbb{N}} \left| \sum_{m=0}^j \sum_{n=0}^k y_{mn} \right| \\ &= \sup_{j,k \in \mathbb{N}} |y_{jk}| = \|y\|_\infty < \infty, \end{aligned}$$

$x = (x_{jk})$ defined by (2.6) is in the space $\mathcal{M}_u(\Delta)$. Hence, T is surjective and norm preserving. This completes the proof of the theorem. $\square$

Now, we give some inclusion relations between the double difference sequence spaces.

Theorem 2.3. *$\mathcal{M}_u$ is the subspace of the space $\mathcal{M}_u(\Delta)$.*

Proof. Let us take $x = (x_{mn}) \in \mathcal{M}_u$. Then, there exists an K such that

$$\sup_{m,n \in \mathbb{N}} |x_{mn}| \leq K$$

for all $m, n \in \mathbb{N}$, one can observe that

$$(2.7) \quad \begin{aligned} |(\Delta x)_{mn}| &= |x_{mn} - x_{m,n-1} - x_{m-1,n} + x_{m-1,n-1}| \\ &\leq |x_{mn}| + |x_{m,n-1}| + |x_{m-1,n}| + |x_{m-1,n-1}|. \end{aligned}$$

Then, we see by taking supremum over $m, n \in \mathbb{N}$ in (2.7) that $\|x\|_\infty \leq 4K$, i.e., $x \in \mathcal{M}_u(\Delta)$.

Now, we see that the inclusion is strict. Let $x = (x_{mn})$ be defined by

$$x_{mn} = mn$$

for all $m, n \in \mathbb{N}$. Then the sequence is in $x \in \mathcal{M}_u(\Delta) \setminus \mathcal{M}_u$. This completes the proof of the theorem. $\square$

Lemma 2.4. [10, Theorem 1.2] $\mathcal{L}_u \subset \mathcal{BS} \subset \mathcal{M}_u$ strictly hold.

Lemma 2.5. [10, Theorem 2.9] Let $v \in \{p, bp, r\}$. Then, the inclusion $\mathcal{BV} \subset \mathcal{C}_v$ and $\mathcal{BV} \subset \mathcal{M}_u$ strictly hold.

Combining Lemma 2.4, Lemma 2.5 and Theorem 2.3, we get the following corollaries.

Corollary 2.6. *The inclusion $\mathcal{L}_u \subset \mathcal{BS} \subset \mathcal{M}_u(\Delta)$ strictly hold.*

Corollary 2.7. *The inclusion $\mathcal{BV} \subset \mathcal{M}_u(\Delta)$ strictly holds.*

Theorem 2.8. *The inclusion $\mathcal{L}_q \subset \mathcal{L}_q(\Delta)$ strictly holds; where $1 \leq q < \infty$.*

Proof. To prove the validity of the inclusion $\mathcal{L}_q \subset \mathcal{L}_q(\Delta)$ for $1 \leq q < \infty$, it suffices to show the existence of a number $K > 0$ such that

$$\|x\|_{\mathcal{L}_q(\Delta)} \leq K\|x\|_{\mathcal{L}_q}$$

for every $x \in \mathcal{L}_q$. Let $x \in \mathcal{L}_q$ and $1 \leq q < \infty$. Then, we obtain

$$\begin{aligned} \|x\|_{\mathcal{L}_q(\Delta)} &= \left\{ \sum_{m,n} |x_{mn} - x_{m,n-1} - x_{m-1,n} + x_{m-1,n-1}|^q \right\}^{1/q} \\ &\leq 4\|x\|_{\mathcal{L}_q}. \end{aligned}$$

This shows that the inclusion $\mathcal{L}_q \subset \mathcal{L}_q(\Delta)$ holds.

Additionally, since the sequence $x = (x_{mn})$ defined by

$$x_{mn} := \begin{cases} 1 & , \quad n = 0 \\ 0 & , \quad \text{otherwise} \end{cases}$$

for all $m, n \in \mathbb{N}$ is in $\mathcal{L}_q(\Delta)$ but not in $\mathcal{L}_q$, as asserted. This completes the proof. $\square$

Theorem 2.9. *The following statements hold:*

- (i) $\mathcal{C}_p$ is the subspace of the space $\mathcal{C}_p(\Delta)$.
- (ii) $\mathcal{C}_{0p}$ is the subspace of the space $\mathcal{C}_{0p}(\Delta)$.
- (iii) $\mathcal{C}_{bp}$ is the subspace of the space $\mathcal{C}_{bp}(\Delta)$.
- (iv) $\mathcal{C}_r$ is the subspace of the space $\mathcal{C}_r(\Delta)$.

Proof. We only prove that the inclusion $\mathcal{C}_p \subset \mathcal{C}_p(\Delta)$ holds. Let us take $x \in \mathcal{C}_p$. Then, for a given $\varepsilon > 0$, there exists an $n_x(\varepsilon) \in \mathbb{N}$ such that

$$|x_{mn} - l| < \frac{\varepsilon}{4}$$

for all $m, n > n_x(\varepsilon)$. Then,

$$\begin{aligned} |(\Delta x)_{mn}| &= |x_{mn} - x_{m,n-1} - x_{m-1,n} + x_{m-1,n-1}| \\ &\leq |x_{mn} - l| + |x_{m,n-1} - l| + |x_{m-1,n} - l| + |x_{m-1,n-1} - l| \\ &< \frac{\varepsilon}{4} + \frac{\varepsilon}{4} + \frac{\varepsilon}{4} + \frac{\varepsilon}{4} = \varepsilon \end{aligned}$$

for sufficiently large m, n which means that $p - \lim(\Delta x)_{mn} = 0$. Hence, $x \in \mathcal{C}_p(\Delta)$ that is to say that $\mathcal{C}_p \subset \mathcal{C}_p(\Delta)$ holds, as expected.

Now, we see that the inclusion is strict. Let $x = (x_{mn})$ be defined by

$$x_{mn} = (m+1)(n+1)$$

for all $m, n \in \mathbb{N}$. It is easy to see that

$$p - \lim(\Delta x)_{mn} = 1.$$

But

$$\lim_{m,n \rightarrow \infty} (m+1)(n+1)$$

which does not tend to a finite limit. Hence $x \notin \mathcal{C}_p$. This completes the proof. $\square$

Lemma 2.10. [10, Theorem 2.3] $\mathcal{CS}_p$ is the subspace of $\mathcal{C}_p$.

Combining Lemma 2.10 and Theorem 2.9, we get the following corollary.

Corollary 2.11. *The inclusion $\mathcal{CS}_p \subset \mathcal{C}_p(\Delta)$ strictly holds.*

3. THE ALPHA- AND BETA-DUALS OF THE NEW SPACES OF DOUBLE SEQUENCES

In this section, we determine the alpha-dual of the space $\mathcal{M}_u(\Delta)$ and the $\beta(r)$ -dual of the space $\mathcal{C}_r(\Delta)$, and $\beta(\vartheta)$ -dual of the space $\mathcal{C}_\eta(\Delta)$ of double sequences, $\vartheta, \eta \in \{p, bp, r\}$. Although the alpha-dual of a space of double sequences is unique, its beta-dual may be more than one with respect to ϑ -convergence.

Theorem 3.1. $\{\mathcal{M}_u(\Delta)\}^\alpha = \mathcal{L}_u$.

Proof. Let $x = (x_{kl}) \in \mathcal{M}_u(\Delta)$ and $z = (z_{kl}) \in \mathcal{L}_u$. Hence, there is a sequence $y = (y_{ij}) \in \mathcal{M}_u$ related with $x = (x_{kl})$ from Theorem 2.2 and there is a positive real number K such that $|y_{ij}| \leq \frac{K}{(k+1)(l+1)}$ for all $i, j \in \mathbb{N}$. So we use the relation (2.6) we have that,

$$\sum_{k,l} |z_{kl}x_{kl}| = \sum_{k,l} \left| z_{kl} \sum_{i=0}^k \sum_{j=0}^l y_{ij} \right| \leq K \sum_{k,l} |z_{kl}| < \infty$$

so $z \in \{\mathcal{M}_u(\Delta)\}^\alpha$, that is

$$(3.1) \quad \mathcal{L}_u \subset \{\mathcal{M}_u(\Delta)\}^\alpha.$$

Conversely, suppose that $z = (z_{kl}) \in \{\mathcal{M}_u(\Delta)\}^\alpha$, that is $\sum_{k,l} |z_{kl}x_{kl}| < \infty$ for all $x = (x_{kl}) \in \mathcal{M}_u(\Delta)$. If $z = (z_{kl}) \notin \mathcal{L}_u$, then $\sum_{k,l} |z_{kl}| = \infty$. Further, if we choose $y = (y_{kl})$ such that

$$y_{kl} := \begin{cases} \frac{1}{(k+1)(l+1)} & , \quad 0 \leq i \leq k, \quad 0 \leq j \leq l \\ 0 & , \quad \text{otherwise} \end{cases}$$

for all $k, l \in \mathbb{N}$. Then, $y \in \mathcal{M}_u$ but

$$\sum_{k,l} |z_{kl}x_{kl}| = \sum_{k,l} \left| z_{kl} \sum_{i=0}^k \sum_{j=0}^l \frac{1}{(k+1)(l+1)} \right| = \sum_{k,l} |z_{kl}| = \infty.$$

Hence, $z \notin \{\mathcal{M}_u(\Delta)\}^\alpha$, this is a contradiction. So, we have the following inclusion,

$$(3.2) \quad \{\mathcal{M}_u(\Delta)\}^\alpha \subset \mathcal{L}_u.$$

Hence, from the inclusions (3.1) and (3.2) we get

$$\{\mathcal{M}_u(\Delta)\}^\alpha = \mathcal{L}_u.$$

□

Now, we may give the β -duals of the spaces with respect to the ϑ -convergence using the technique in [12] and [13] for the single sequences.

The conditions for a 4-dimensional matrix to transform the spaces $\mathcal{C}_{bp}, \mathcal{C}_r$ and $\mathcal{C}_p$ into the space $\mathcal{C}_{bp}$ are well known (see for example [14, 15]).

Lemma 3.2. The matrix $A = (a_{mnij})$ is in $(\mathcal{C}_r : \mathcal{C}_\vartheta)$ if and only if the following conditions hold:

$$(3.3) \quad \sup_{m,n \in \mathbb{N}} \sum_{i,j} |a_{mnij}| < \infty,$$

$$(3.4) \quad \exists v \in \mathbb{C} \ni \vartheta - \lim_{m,n \rightarrow \infty} \sum_{i,j} a_{mnij} = v,$$

$$(3.5) \quad \exists (a_{ij}) \in \Omega \ni \vartheta - \lim_{m,n \rightarrow \infty} a_{mnij} = a_{ij} \quad \text{for all } i, j \in \mathbb{N},$$

$$(3.6) \quad \exists u^{j_0} \in \mathbb{C} \ni \vartheta - \lim_{m,n \rightarrow \infty} \sum_i a_{mnij_0} = u^{j_0} \quad \text{for fixed } j_0 \in \mathbb{N},$$

$$(3.7) \quad \exists v_{i_0} \in \mathbb{C} \ni \vartheta - \lim_{m,n \rightarrow \infty} \sum_j a_{mnij_0} = v_{i_0} \quad \text{for fixed } i_0 \in \mathbb{N}.$$

Lemma 3.3. The matrix $A = (a_{mni j})$ is in $(\mathcal{C}_{bp} : \mathcal{C}_\vartheta)$ if and only if the conditions (3.3)-(3.5) of Lemma 3.2 hold, and

$$(3.8) \quad \vartheta - \lim_{m,n \rightarrow \infty} \sum_i |a_{mni j_0} - a_{i j_0}| = 0 \text{ for each fixed } j_0 \in \mathbb{N},$$

$$(3.9) \quad \vartheta - \lim_{m,n \rightarrow \infty} \sum_j |a_{mni_0 j} - a_{i_0 j}| = 0 \text{ for each fixed } i_0 \in \mathbb{N}.$$

Lemma 3.4. The matrix $A = (a_{mni j})$ is in $(\mathcal{C}_p : \mathcal{C}_\vartheta)$ if and only if the conditions (3.3)-(3.5) of Lemma 3.2 hold, and

$$(3.10) \quad \forall i \in \mathbb{N} \exists J \in \mathbb{N} \ni a_{mni j} = 0 \text{ for } j > J \text{ for all } m, n \in \mathbb{N},$$

$$(3.11) \quad \forall j \in \mathbb{N} \exists I \in \mathbb{N} \ni a_{mni j} = 0 \text{ for } i > I \text{ for all } m, n \in \mathbb{N}.$$

Theorem 3.5. Define the sets

$$\begin{aligned} F_1 &= \left\{ a = (a_{ij}) \in \Omega : \sum_{i,j} (i+1)(j+1)|a_{ij}| < \infty \right\}, \\ F_2 &= \left\{ a = (a_{ij}) \in \Omega : r - \lim_{m,n \rightarrow \infty} \sum_i^m \sum_{p=i}^m \sum_{q=j_0}^n a_{pq} \text{ exists for each fixed } j_0 \right\}, \\ F_3 &= \left\{ a = (a_{ij}) \in \Omega : r - \lim_{m,n \rightarrow \infty} \sum_j^m \sum_{p=i_0}^m \sum_{q=j}^n a_{pq} \text{ exists for each fixed } i_0 \right\}. \end{aligned}$$

Then, $\{\mathcal{C}_r(\Delta)\}^{\beta(r)} = F_1 \cap F_2 \cap F_3$.

Proof. Let $x = (x_{ij}) \in \mathcal{C}_r(\Delta)$. Then, there exists a sequence $y = (y_{mn}) \in \mathcal{C}_r$. Consider the following equality

$$\begin{aligned} z_{mn} &= \sum_{i=0}^m \sum_{j=0}^n a_{ij} x_{ij} = \sum_{i=0}^m \sum_{j=0}^n \left(\sum_{p=0}^i \sum_{q=0}^j y_{pq} \right) a_{ij} \\ &= \sum_{i=0}^m \sum_{j=0}^n \left(\sum_{p=i}^m \sum_{q=j}^n a_{pq} \right) y_{ij} \\ &= \sum_{i=0}^m \sum_{j=0}^n b_{mni j} y_{ij} = (By)_{ij} \end{aligned}$$

for all $m, n \in \mathbb{N}$. Hence we can define the four-dimensional matrix $B = (b_{mni j})$ as following

$$(3.12) \quad b_{mni j} := \begin{cases} \sum_{p=i}^m \sum_{q=j}^n a_{pq} & , \quad 0 \leq i \leq m, \quad 0 \leq j \leq n, \\ 0 & , \quad \text{otherwise.} \end{cases}$$

Thus we see that $ax = (a_{mn} x_{mn}) \in \mathcal{CS}_r$ whenever $x = (x_{mn}) \in \mathcal{C}_r(\Delta)$ if and only if $z = (z_{mn}) \in \mathcal{C}_r$ whenever $y = (y_{mn}) \in \mathcal{C}_r$. This means that $a = (a_{mn}) \in \{\mathcal{C}_r(\Delta)\}^{\beta(r)}$ if and only if $B \in (\mathcal{C}_r : \mathcal{C}_r)$. Therefore, we consider the following equality and equation

$$\begin{aligned} \sup_{m,n \in \mathbb{N}} \sum_{i=0}^m \sum_{j=0}^n |b_{mni j}| &\leq \sup_{m,n \in \mathbb{N}} \sum_{i=0}^m \sum_{j=0}^n \left(\sum_{p=i}^m \sum_{q=j}^n |a_{pq}| \right) \\ &= \sup_{m,n \in \mathbb{N}} \sum_{i=0}^m \sum_{j=0}^n \left(\sum_{p=0}^i \sum_{q=0}^j |a_{ij}| \right) \\ &= \sup_{m,n \in \mathbb{N}} \sum_{i=0}^m \sum_{j=0}^n (i+1)(j+1)|a_{ij}|, \end{aligned} \tag{3.13}$$

$$\begin{aligned}
(3.14) \quad r - \lim_{m,n \rightarrow \infty} \sum_{i,j} b_{mnij} &= r - \lim_{m,n \rightarrow \infty} \sum_{i=0}^m \sum_{j=0}^n \left(\sum_{p=0}^i \sum_{q=0}^j a_{pq} \right) \\
&= \sum_{i,j} \left(\sum_{p=0}^i \sum_{q=0}^j a_{pq} \right).
\end{aligned}$$

Then, we derive from the condition (3.3)-(3.5) that

$$(3.15) \quad \sum_{i,j} (i+1)(j+1)|a_{ij}| < \infty.$$

Further, from Lemma 3.2 conditions (3.6) and (3.7),

$$(3.16) \quad r - \lim_{m,n \rightarrow \infty} \sum_i b_{mnij_0} = r - \lim_{m,n \rightarrow \infty} \sum_i \sum_{p=i}^m \sum_{q=j_0}^n a_{pq}$$

exists for each fixed $j_0 \in \mathbb{N}$ and

$$(3.17) \quad r - \lim_{m,n \rightarrow \infty} \sum_j b_{mnij_0} = r - \lim_{m,n \rightarrow \infty} \sum_i \sum_{p=i_0}^m \sum_{q=j}^n a_{pq}$$

exists for each fixed $i_0 \in \mathbb{N}$. This show that $\{\mathcal{C}_r(\Delta)\}^{\beta(r)} = F_1 \cap F_2 \cap F_3$ which completes the proof. $\square$

Now, we may give our theorem exhibiting the $\beta(\vartheta)$ -dual of the series space $\mathcal{C}_\eta(\Delta)$ in the case $\eta, \vartheta \in \{p, bp, r\}$, without proof.

Theorem 3.6. $\{\mathcal{C}_\eta(\Delta)\}^{\beta(\vartheta)} = \{a = (a_{mn}) \in \Omega : B = (b_{mnij}) \in (\mathcal{C}_\eta : \mathcal{C}_\vartheta)\}$ where $B = (b_{mnij})$ is defined by (3.12).

4. CHARACTERIZATION OF SOME CLASSES OF FOUR DIMENSIONAL MATRICES

In the present section, we characterize the matrix transformations from the space $\mathcal{C}_r(\Delta)$ to the double sequence space $\mathcal{C}_\vartheta$. Although the theorem characterizing the class $(\mu : \mathcal{C}_\vartheta(\Delta))$ are stated and proved, the necessary and sufficient conditions on a four dimensional matrix belonging to the classes $(\mathcal{C}_r : \mathcal{C}_r(\Delta))$ and $(\mathcal{C}_{bp} : \mathcal{C}_{bp}(\Delta))$ also given without proof.

Theorem 4.1. $A = (a_{mnkl}) \in (\mathcal{C}_r(\Delta) : \mathcal{C}_\vartheta)$ if and only if the following conditions hold:

$$(4.1) \quad \sup_{m,n} \sum_{k,l} \left| \sum_{p=k}^{\infty} \sum_{q=l}^{\infty} a_{mnpq} \right| < \infty,$$

$$(4.2) \quad \vartheta - \lim_{s,t \rightarrow \infty} \sum_{i=0}^s \sum_{p=i}^s \sum_{p=j_0}^t a_{mnpq} - \text{exists for fixed } j_0,$$

$$(4.3) \quad \vartheta - \lim_{s,t \rightarrow \infty} \sum_{j=0}^t \sum_{p=i_0}^s \sum_{p=j}^t a_{mnpq} - \text{exists for fixed } i_0,$$

$$(4.4) \quad \vartheta - \lim_{m,n} \sum_{p=k}^{\infty} \sum_{q=l}^{\infty} a_{mnpq} = a_{kl} \text{ for all } k, l \in \mathbb{N},$$

$$(4.5) \quad \exists u^{l_0} \in \mathbb{C} \ni \vartheta - \lim_{m,n} \sum_k \sum_{p=k}^{\infty} \sum_{q=l_0}^{\infty} a_{mnpq} = u^{l_0} \text{ for fixed } l_0 \in \mathbb{N},$$

$$(4.6) \quad \exists v_{k_0} \in \mathbb{C} \ni \vartheta - \lim_{m,n} \sum_l \sum_{p=k_0}^{\infty} \sum_{q=l}^{\infty} a_{mnpq} = v_{k_0} \text{ for fixed } k_0 \in \mathbb{N},$$

$$(4.7) \quad \exists v \in \mathbb{C} \ni \vartheta - \lim_{m,n} \sum_{k,l} \sum_{p=k}^{\infty} \sum_{q=l}^{\infty} a_{mnpq} = v.$$

Proof. Let us take any $x = (x_{mn}) \in \mathcal{C}_r(\Delta)$ and define the sequence $y = (y_{kl})$ by

$$y_{kl} = x_{kl} - x_{k-1,l} - x_{k,l-1} + x_{k-1,l-1}; \quad (k, l \in \mathbb{N}).$$

Then, $y = (y_{kl}) \in \mathcal{C}_r$ by Theorem 2.2. Now, for the (s, t) th rectangular partial sum of the series $\sum_{j,k} a_{mnjk} x_{jk}$, we derive that

$$\begin{aligned} (Ax)_{mn}^{[s,t]} &= \sum_{j=0}^s \sum_{k=0}^t a_{mnjk} x_{jk} \\ &= \sum_{j=0}^s \sum_{k=0}^t \left(\sum_{p=0}^j \sum_{q=0}^k y_{pq} \right) a_{mnjk} \\ (4.8) \quad &= \sum_{j=0}^s \sum_{k=0}^t \left(\sum_{p=j}^s \sum_{q=k}^t a_{mnpq} \right) y_{jk} \end{aligned}$$

for all $m, n, s, t \in \mathbb{N}$. Define the matrix $B_{mn} = (b_{mnjk}^{[s,t]})$ by

$$(4.9) \quad b_{mnjk}^{[s,t]} := \begin{cases} \sum_{p=j}^s \sum_{q=k}^t a_{mnpq} & , \quad 0 \leq j \leq s, \quad 0 \leq k \leq t, \\ 0 & , \quad \text{otherwise.} \end{cases}$$

Then, the equality (4.8) may be rewritten as

$$(4.10) \quad (Ax)_{mn}^{[s,t]} = (B_{mn}y)_{[s,t]}.$$

Then, the convergence of the rectangular partial sums $(Ax)_{mn}^{[s,t]}$ in the regular sense for all $m, n \in \mathbb{N}$ and for all $x \in \mathcal{C}_r(\Delta)$ is equivalent of saying that $B_{mn} \in (\mathcal{C}_r : \mathcal{C}_\vartheta)$. Hence, the following conditions

$$(4.11) \quad \sum_{k,l} (k+1)(l+1) |a_{mnkl}| < \infty,$$

$$(4.12) \quad \vartheta - \lim_{s,t \rightarrow \infty} \sum_{i=0}^s \sum_{p=i}^s \sum_{p=j_0}^t a_{mnpq} - \text{exists for fixed } j_0,$$

$$(4.13) \quad \vartheta - \lim_{s,t \rightarrow \infty} \sum_{j=0}^t \sum_{p=i_0}^s \sum_{p=j}^t a_{mnpq} - \text{exists for fixed } i_0$$

must be satisfied for every fixed $m, n \in \mathbb{N}$. In this case,

$$\vartheta - \lim_{s,t \rightarrow \infty} b_{mnjk}^{[s,t]} = \sum_{p=j}^{\infty} \sum_{q=k}^{\infty} a_{mnpq},$$

$$\vartheta - (Ax)_{mn}^{[s,t]} = r - \lim (B_{mn}y)$$

hold. Thus, we derive from the two-sided implication " Ax is in C_r whenever $x \in C_r(\Delta)$ if and only if $B = (\sum_{p=j}^{\infty} \sum_{q=k}^{\infty} a_{mnpq})_{mn} \in (C_r : C_{\vartheta})$ ", we have Lemma 3.2 that

$$(4.14) \quad \sup_{m,n} \sum_{k,l} \left| \sum_{p=k}^{\infty} \sum_{q=l}^{\infty} a_{mnpq} \right| < \infty,$$

$$(4.15) \quad \vartheta - \lim_{m,n} \sum_{p=k}^{\infty} \sum_{q=l}^{\infty} a_{mnpq} = a_{kl} \text{ for all } k, l \in \mathbb{N},$$

$$(4.16) \quad \exists u^{l_0} \in \mathbb{C} \ni \vartheta - \lim_{m,n} \sum_k \sum_{p=k}^{\infty} \sum_{q=l_0}^{\infty} a_{mnpq} = u^{l_0} \text{ for fixed } l_0 \in \mathbb{N},$$

$$(4.17) \quad \exists v_{k_0} \in \mathbb{C} \ni \vartheta - \lim_{m,n} \sum_l \sum_{p=k_0}^{\infty} \sum_{q=l}^{\infty} a_{mnpq} = v_{k_0} \text{ for fixed } k_0 \in \mathbb{N},$$

$$(4.18) \quad \exists v \in \mathbb{C} \ni \vartheta - \lim_{m,n} \sum_{k,l} \sum_{p=k}^{\infty} \sum_{q=l}^{\infty} a_{mnpq} = v.$$

Now, from the conditions (4.11)-(4.18), we have that $A = (a_{mnkl}) \in (C_r(\Delta) : C_v)$ if and only if the conditions (4.1)-(4.7) hold. This completes the proof. $\square$

Theorem 4.2. Suppose that the elements of the four dimensional infinite matrices $E = (e_{mnkl})$ and $F = (f_{mnkl})$ are connected with the relation

$$(4.19) \quad f_{mnkl} = \sum_{i=m-1}^m \sum_{j=n-1}^n (-1)^{m+n-i-j} e_{ijkl}$$

for all $k, l, m, n \in \mathbb{N}$ and μ be any given space of double sequences. Then, $E \in (\mu : C_{\vartheta}(\Delta))$ if and only if $F \in (\mu : C_{\vartheta})$.

Proof. Let $x = (x_{kl}) \in \mu$ and consider the following equality with (4.19)

$$(4.20) \quad \sum_{i=m-1}^m \sum_{j=n-1}^n \sum_{k=s-1}^s \sum_{l=t-1}^t (-1)^{m+n-i-j} e_{ijkl} x_{kl} = \sum_{k=s-1}^s \sum_{l=t-1}^t f_{mnkl} x_{kl}$$

for all $m, n, s, t \in \mathbb{N}$. By letting $s, t \rightarrow \infty$ in (4.20) one can derive that

$$(4.21) \quad \sum_{i=m-1}^m \sum_{j=n-1}^n (-1)^{m+n-i-j} (Ex)_{ij} = (Fx)_{mn} \text{ for all } m, n \in \mathbb{N}.$$

Therefore, it is seen by (4.21) that $Ex \in C_{\vartheta}(\Delta)$ if and only if $Fx \in C_{\vartheta}$ whenever $x \in \mu$. This step completes the proof. $\square$

Of course, Theorem 4.2 has several consequences depending on the choice of the sequence space μ . Prior to giving some results as an application of this idea, we need the following lemmas:

Lemma 4.3. [14, 16, 17] $A = (a_{mnkl}) \in (C_r : C_r)$ if and only if

$$(4.22) \quad \sup_{m,n \in \mathbb{N}} \sum_{k,l} |a_{mnkl}| < \infty,$$

$$(4.23) \quad \exists (a_{kl}) \in \Omega \ni r - \lim_{m,n \rightarrow \infty} a_{mnkl} = a_{kl} \text{ for each } k, l \in \mathbb{N},$$

$$(4.24) \quad \exists v \in \mathbb{C} \ni r - \lim_{m,n \rightarrow \infty} \sum_{k,l} a_{mnkl} = v,$$

$$(4.25) \quad \exists u^{l_0}, v_{k_0} \in \mathbb{C} \ni r - \lim_{m,n \rightarrow \infty} \sum_k a_{mnkl_0} = u^{l_0} \text{ and}$$

$$r - \lim_{m,n \rightarrow \infty} \sum_l a_{mnk_0l} = v_{k_0} \text{ for any } k_0, l_0 \in \mathbb{N}.$$

Lemma 4.4. [14, 16, 17] $A = (a_{mnkl}) \in (\mathcal{C}_{bp} : \mathcal{C}_{bp})$ if and only if

$$(4.26) \quad \sup_{m,n \in \mathbb{N}} \sum_{k,l} |a_{mnkl}| < \infty,$$

$$(4.27) \quad \exists (a_{kl}) \in \Omega \ni bp - \lim_{m,n \rightarrow \infty} a_{mnkl} = a_{kl} \quad \text{for each } k, l \in \mathbb{N},$$

$$(4.28) \quad \exists v \in \mathbb{C} \ni bp - \lim_{m,n \rightarrow \infty} \sum_{k,l} a_{mnkl} = v,$$

$$(4.29) \quad bp - \lim_{m,n \rightarrow \infty} \sum_k |a_{mnkl_0} - a_{kl_0}| = 0 \quad \text{and}$$

$$bp - \lim_{m,n \rightarrow \infty} \sum_l |a_{mnk_0l} - a_{k_0l}| = 0 \quad \text{for any } k_0, l_0 \in \mathbb{N}.$$

Corollary 4.5. Suppose that the relation (4.19) holds between the elements of the four dimensional infinite matrices $E = (e_{mnkl})$ and $F = (f_{mnkl})$. Then, the following statements hold:

(i) $E = (e_{mnjk}) \in (\mathcal{C}_r : \mathcal{C}_r(\Delta))$ if and only if the conditions (4.22)-(4.25) hold with f_{mnkl} instead of a_{mnkl} .

(ii) $E = (e_{mnjk}) \in (\mathcal{C}_{bp} : \mathcal{C}_{bp}(\Delta))$ if and only if the conditions (4.26)-(4.29) hold with f_{mnkl} instead of a_{mnkl} .

ACKNOWLEDGEMENT

We wish to thank the referees for their valuable suggestions and comments which improved the paper considerably.

REFERENCES

- [1] J. Boss, Classical and Modern Methods in Summability, Oxford University Press, Newyork, 2000.
- [2] F. Mòricz, Extensions of the spaces c and c_0 from single to double sequences, Acta Math. Hungar., **57** (1991), 129-136.
- [3] M. Zeltser, Investigation of double sequence spaces by soft and hard analitic methods, Dissertationes Mathematicae Universtatis Tartuenss **25**, Tartu University Press, Univ. of Tartu, Faculty of Mathematics and Computer Science, Tartu, 2001.
- [4] M. Zeltser, On conservative matrix methods for double sequence spaces, Acta Math. Hung., **95**(3) (2002), 225-242.
- [5] A. Gökhan and R. Çolak, The double sequence spaces $c_2^P(p)$ and $c_2^{BP}(p)$, Appl. Math. Comput., **157**(2) (2004), 491-501.
- [6] A. Gökhan and R. Çolak, Double sequence space $\ell_2^\infty(p)$, Appl. Math. Comput., **160** (2005), 147-153.
- [7] M. Mursaleen and O.H.H. Edely, Statistical convergence of double sequences, J. Math. Anal. Appl., **288**(1) (2003), 223-231.
- [8] M. Mursaleen, Almost strongly regular matrices and a core theorem for double sequences, J. Math. Anal. Appl., **293**(2) (2004), 523-531.
- [9] M. Mursaleen and O.H.H. Edely, Almost convergence and a core theorem for double sequences, J. Math. Anal. Appl., **293**(2) (2004), 532-540.
- [10] B. Altay and F. Başar, Some new spaces of double sequences, J. Math. Anal. Appl., **309**(1) (2005), 70-90.
- [11] F. Başar and Y. Sever, The space $\mathcal{L}_q$ of double sequences, Math. J. Okayama Univ., **51** (2009), 149-157.
- [12] B. Altay and F. Başar, Matrix mappings on the space $bs(p)$ and its α -, β - and γ -duals, Aligarh Bull. Math., **21** (2002), 79-91.
- [13] B. Altay and F. Başar, On the space of sequences of p -bounded variation and related matrix mappings, Ukrainian Math. J., **55**(1) (2003), 136-147.
- [14] H. J. Hamilton, Transformations of multiple sequences, Duke Math. J., **2** (1936), 29-60.
- [15] M. Zeltser, M. Mursaleen and S. A. Mohiuddine, On almost conservative matrix methods for double sequence spaces, Publ. Math. Debrecen, **75** (2009), 387-399.
- [16] G. M. Robison, Divergent double sequences and series, Amer. Math. Soc. Trans., **28** (1926), 50-73.
- [17] M. Zeltser, On conservative matrix methods for double sequences, Acta Math. Hung., **95**(3) (2002), 225-242.

Current address: Gaziosmanpaşa University, Faculty of Arts and Science, Department of Mathematics, Tokat-60240/TURKEY
E-mail address: `serkandemiriz@gmail.com`, `osman-duyar@hotmail.com`